

# A study of the scenario with nearly degenerate Majorana neutrinos

Francesco Vissani

*International Centre for Theoretical Physics,  
Strada Costiera 11, I-34013 Trieste, Italy*

(August 28, 1997)

Motivated by cosmological considerations, and by the atmospheric and solar neutrino flux deficits, we consider the scenario in which the three Standard Model neutrinos are endowed with a nearly equal Majorana mass in the eV range. Combined constraints coming from (1) direct search for electron neutrino mass, (2) absence of neutrinoless double beta decay, and (3) unsuppressed electron neutrino flux in present reactor experiments, imply a quite well specified pattern of neutrino masses and mixing angles. We discuss the experimental tests of the model and comment on the features of this scenario.

## I. NEARLY DEGENERATE MAJORANA NEUTRINOS

### A. Phenomenological character of the approach

The possibility that the neutrinos  $\nu_e$ ,  $\nu_\mu$  and  $\nu_\tau$  produced in weak interactions are coherent combinations of almost degenerate mass eigenstates has attracted great attention in the particle physics community [1–12]. This is motivated by the fact that the mass splitting required to explain the atmospheric and the solar neutrino flux deficits in terms of neutrino oscillations are much smaller than the mass required for cosmological purposes, as will be discussed below in this first section.

Most of the works focused on the *ansatze* for the neutrino mass matrix:  $\mathcal{M} = m_0 \times \mathbb{1} + \text{small corrections}$ , and on the study of the corresponding theoretical motivations and phenomenological implications. Inevitably, this *ansatze* implies a signal for neutrinoless double beta decay ( $0\nu\beta\beta$ ) experiments if  $m_0 = \text{few eV's}$ , of interest for cosmological considerations. Allowing a departure from this mass pattern, the authors of [9] outlined a scenario in which the neutrinos have masses in the sub-eV range and could produce a signal in next generation searches of  $0\nu\beta\beta$ . It was later proven that it is possible [11] to incorporate a massive neutrino in the eV range in the context of models with abelian family symmetry, in a way marginally consistent with the  $0\nu\beta\beta$  bounds.

In the present work we do not commit *a priori* ourselves to any given pattern for the neutrino mass matrix, and concentrate on finding a picture of the neutrino masses and mixing angles that is consistent with the phenomenological observations, and with the hypothesis of massive neutrinos in the cosmologically interesting range. We will show that these hypotheses are quite restrictive, implying that the corresponding scenario is likely to be confirmed or disproved by forthcoming experiments.

In its phenomenological character, our approach is very similar to the one in [12]. We emphasize in the present study the importance of the direct search for neutrino masses, as crucial test of the scenario under discussion. Another difference with [12] is in the fact that the authors of that work concentrate on the MSW solution of the solar neutrino problem [13–15], whereas we focus on the vacuum oscillation solution [16]; in this respect, the two analysis complement each other.

### B. Definition of the scenario

The lagrangian of neutrino masses in the basis where the charged lepton masses are diagonal is (bispinorial notation):

$$\mathcal{L}_{\text{mass}} = - \sum_{\ell \text{ and } \ell' = e, \mu, \tau} \mathcal{M}_{\ell\ell'} N_\ell(x) N_{\ell'}(x) + \text{h.c.}; \quad (1)$$

the Lorentz indices are contracted according to  $N^A N_A$ ,  $A = 1, 2$ . The symmetric matrix  $\mathcal{M}$  can be decomposed as:

$$\mathcal{M}_{\ell\ell'} = \sum_{i=1,2,3} U_{\ell i}^* U_{\ell' i} m_{\nu_i} \quad (2)$$

(in compact notation:  $\mathcal{M} = U^* m U^\dagger$ ), where  $U$  is the unitary matrix of mixing angles and phases. The three nonnegative and nondecreasing real numbers  $m_{\nu_i}$  are the Majorana neutrino masses. Introducing the bispinors  $N_i(x)$ :

$$N_\ell(x) = \sum_{i=1,2,3} U_{\ell i} N_i(x) \quad \ell = e, \mu, \tau \quad (3)$$

and the Majorana four-spinor in the chiral representation:  $\nu_{i\alpha}(x) = (N_{iA}(x), \bar{N}_i^{\dot{A}}(x))$  we can finally rewrite the mass lagrangian in the standard form:

$$\mathcal{L}_{\text{mass}} = -\frac{1}{2} \sum_{i=1,2,3} m_{\nu_i} \bar{\nu}_i(x) \nu_i(x). \quad (4)$$

The mass difference is denoted as:

$$\Delta m_{ij}^2 = m_{\nu_i}^2 - m_{\nu_j}^2. \quad (5)$$

It enters for instance in the expression  $P_{\ell\ell'} = |\sum_i U_{\ell i}^* U_{\ell' i} \exp(-iE_i t)|^2$  of the probability of oscillation in vacuum from a flavour  $\ell$  to  $\ell'$ . In fact we can replace:  $\exp(-iE_i t)/\exp(-iE_j t)$  by  $\exp(-i\Delta m_{ij}^2 t/(2p))$ , under the assumption (which holds in the cases of interest) that the neutrino momentum  $p$  is much larger than the neutrino masses.

The scenario under discussion can now be defined by letting in first approximation:

$$m_{\nu_i} = m_0 \quad i = 1, 2, 3. \quad (6)$$

The value of  $m_0$  is fixed below, with the help of cosmological considerations. Before passing to that, we further specify two sub-cases of this scenario, by making the assumption that the squared mass differences are in ranges of interest to the solar and the atmospheric neutrino problems. Case A is defined by:

$$\Delta m_{21}^2 = \Delta m_\odot^2 \approx 10^{-10} \text{eV}^2, \quad \Delta m_{32}^2 = \Delta m_{\text{atm}}^2 \approx 10^{-2} \text{eV}^2; \quad (7)$$

in case B the two mass differences are inverted (compare with fig. 1 in last section). Since, apart from this fact (and neglecting matter effects on the propagation of neutrinos), the two cases are phenomenologically equivalent, we will concentrate the discussion on case A.

### C. Neutrino masses of cosmological interest

There are various indications of the existence of dark matter (DM—see for instance [17]), that may constitute the main component of the Universe. Once the amount of dark matter, ordinary matter and, possible other components are specified, one can attempt a comparison of the observed structure with the one predicted in terms of the spectrum of the primordial fluctuations. The gross features of the Universe can be explained by the cold DM model, that assumes (1) that the dominant component of DM is cold (non relativistic at the moment of decoupling), (2) a nearly scale invariant (Harrison-Zel'dovich) primordial spectrum, (3) and that the parameter  $\Omega$  is 1, which implies that the total energy density is equal to the critical density  $\rho_c$  (the latter two hypothesis are suggested by inflationary scenarios). However COBE data have indicated the need of a certain departure from this simple cosmological model [19]. The hot+cold DM model [19,20] achieves an optimal agreement with the data if the hot DM constitutes the 20 – 30% of the energy density, the cold one the 70 – 60%, and the baryonic matter approximatively 1 tenth of the total energy density<sup>1</sup>.

Massive neutrinos are natural candidates for hot DM, and this is the assumption we are interested to consider. The number density of neutrinos  $n_\nu$  can be computed in the Big-Bang model [24]. Once they become non-relativistic, their energy density reduces simply to  $\rho_\nu = \sum_i m_{\nu_i} n_\nu$ . Assuming  $\Omega = 1$ , and introducing the ratio of  $\rho_\nu$  to the critical density  $\rho_c$  (proportional to  $H_0^2$ ) one arrives at the expression that relates the neutrino masses to the cosmological parameters:

$$\sum_i m_{\nu_i} = 6.6 \text{ eV} \times \left[ \frac{\rho_\nu}{0.2 \rho_c} \right] \times \left[ \frac{h}{0.6} \right]^2. \quad (8)$$

---

<sup>1</sup>In this model, the favoured value of the Hubble constant  $H_0 = h \times 100 \text{ km sec}^{-1} \text{ Mpc}^{-1}$  turns out to be around  $h = 0.5$ . In [21] the observational range  $h = 0.65 - 0.85$  is suggested; a more recent analysis [22] gives  $h = 0.6 - 0.8$  as the reference range. Even lower values,  $h = 0.5 - 0.7$  have been advocated in [23].

The neutrino mass differences implied by solar and atmospheric neutrino deficits (7) are small in comparison with the mass in eq. (8), and this facts suggest the hypothesis of nearly degenerate neutrinos. Using eqs. (6) and (8) together with the informations from the study of the structure evolution, we are lead to the following reference value for  $m_0 = \sum_i m_{\nu_i}/3$ :

$$m_0 = m(\text{cosm.}) = 2.2 \text{ eV}. \quad (9)$$

This corresponds to  $h = 0.6$  and  $\rho_\nu = 0.2 \rho_c$ , or, with approximation of 5%, to  $h = 0.5$  and  $\rho_\nu = 0.3 \rho_c$ . Progress in the determination of  $H_0$  and  $\Omega$  will provide tests of the hot+cold DM scenario, and also, if the scenario will be confirmed, a better determination of the parameter  $m_0$ .

We briefly comment on the possibility that massive neutrinos may constitute the major component of the DM. It was suggested that this could happen if special seeds of primordial fluctuations like those originated by the cosmic strings [25] play a major role in the process of structure evolution, or, alternatively, if neutrinos participate in exotic long-range (nongravitational) interactions [27]. However, apart from any theoretical justification, if this alternative scenario is correct, and  $\Omega = 1$ , the reference value of the parameter  $m_0$  would be larger than 10 eV, since  $\rho_\nu$  in eq. (8) increases almost 5 times.

Of course, it is a major assumption to consider the absence of light sterile neutrinos, that could play various roles, and in particular could give the dominant contribution to the hot DM content of the Universe [28–30]. The hypothesis of (Standard Model) singlet neutrinos could be tested if they are mixed with the ordinary ones, and participate in the oscillations. Regarding this point, if neutrino masses are the explanation not only of the solar and of the atmospheric neutrino problems, but *also* of the LSND data on  $\bar{\nu}_\mu - \bar{\nu}_e$  [31,32] and on  $\nu_\mu - \nu_e$  flux conversion [33], major modifications of the scenario, like the introduction of a sterile neutrino would be required. Therefore, in order to check the scenario under discussion (in which no light sterile neutrino is present and a negligible signal at LSND is expected) it will be crucial to see if the LSND findings will be confirmed by the forthcoming upgraded KARMEN [34] and/or by other experiments.

## II. CONSTRAINTS FROM PHENOMENOLOGY

In this section we discuss the bounds on the neutrino masses and mixing angles that can be deduced from the absence (1) of positive indications of electron neutrino mass in the tritium decay experiments; (2) of neutrinoless double beta decay signal; (3) of appreciable disappearance of electron neutrino flux in reactor experiments. The constraints are considered in order of relevance for the scenario with three almost degenerate neutrinos<sup>2</sup>.

If one (or more) of the related experiments finds a signal, one (or more) parameters of the model will be determined. We stress that, in the model under discussion, (i) the direct detection of neutrino mass seems to be possible, and that (ii) condition (2) implies a strong restriction on the mixing angles, that will further strengthen if future  $0\nu\beta\beta$  experiments will continue to give a null result.

### A. Direct search for neutrino masses

Let us denote by  $d\Gamma[m_{\nu_i}]/dE_e$  the differential electron spectrum in beta decay supposing that  $\nu_i$  coincides with the electron neutrino. In the general case, in which the electron neutrino is not a mass eigenstate, the beta spectrum  $d\Gamma/dE_e$  is the weighted combination:

$$\frac{d\Gamma}{dE_e} = \sum_{i=1,2,3} \frac{d\Gamma[m_{\nu_i}]}{dE_e} |U_{ei}|^2 \Theta(\Delta M - m_{\nu_i} - E_e), \quad (10)$$

where the Heaveside function  $\Theta(x)$  enters the formula due to kinematical requirements, and  $\Delta M$  is the difference between the mass of the decaying nucleus and that of the final nucleus. In our case, due to (6), the unitarity relation  $\sum_i |U_{ei}|^2 = 1$  implies a constraint straightforwardly on the common mass  $m_0$ .

Stringent limits can be derived by the experimental study of endpoint spectrum in tritium decay, after taking into account in some way the problematic indication of negative masses squared common to most recent experiments

---

<sup>2</sup>The importance of the direct searches for neutrino mass has been not, in our opinion, sufficiently emphasized in the literature, even if the existence of this constraint was realized since the beginning (see for instance fig. 2 in reference [2]).

(corresponding to the presence of a *bump* in the endpoint region). In the most recent experimental analysis [35], still under progress, new parameters unrelated to the neutrino mass are introduced<sup>3</sup>. The upper limit obtained,  $m_\nu^2 = 1.5 \pm 5.9(\text{stat.}) \pm 3.6(\text{syst.}) \text{ eV}^2$ , implies:

$$m_0 < m(^3\text{H}) = 3.9 \text{ eV at 95\% CL}, \quad (11)$$

summing the errors in quadrature. The limit (11) is not very different from those obtained by other experiments; it could be further strengthened to 2.2 eV if the bump would be fully understood [35].

The limits on neutrino mass from SN1987A obtained by the measured spread in time (which is related to the original spread in energy and to the duration of the neutrino pulse) are significantly looser,  $m(\text{SN1987A}) \lesssim 15 \text{ eV}$ . However, the observation of neutrinos from a very distant supernova could yield a competitive or even better bound in future.

The following remarks should clearly illustrate that, in the scenario with nearly degenerate neutrinos, the direct search for neutrino mass is of cosmological relevance:

1. The “preferred” value (9) is quite close to the limit (11):  $m_0 \approx 2 \times m(^3\text{H})$ . Therefore, the hot+cold DM model can be ruled out or confirmed by a slight increase in the precision of the measurements. We stress once more the importance of a clear theoretical description of the structure of the endpoint spectrum in tritium decay studies.
2. For  $\rho_\nu = 0.2 \rho_c$  the limit (11) disfavors  $h > 0.8$  in the hot+cold DM model. Increasingly precise measurements of the cosmological parameters and numerical simulations will be important in narrowing the parameter space of the model.
3. The scenarios in which  $\rho_\nu$  dominates the energy density of Universe and that have  $\Omega = 1$  require  $m_0 \gtrsim 10 \text{ eV}$ , and therefore can already be excluded in the present context.

## B. Neutrinoless double beta decay and large mixing

The limit (11) (the preferred value (8)) are approximatively 6 times (3 times) larger than the limit obtained by searches of neutrinoless double beta decay [37]:

$$|\mathcal{M}_{ee}| = \left| \sum_{i=1,2,3} U_{ei}^2 m_{\nu_i} \right| < m(0\nu\beta\beta) = 0.68 \text{ eV} \quad \text{at 95\% CL}. \quad (12)$$

We disregard in the following the possibility of large errors in the estimation of the hadronic matrix elements. The bound (12) implies a cancellation in the contributions to  $|\mathcal{M}_{ee}|$  with a precision of  $m(0\nu\beta\beta)/m_0 \approx 30 \%$ . Let us first analyze the case in which the neutrinoless double beta decay is strongly suppressed, which implies  $\sum_i U_{ei}^2 \approx 0$ . This equation can be represented geometrically as a triangle in the complex plane. Therefore it can be satisfied only if the sum of the length of two sides is not smaller than the length of the third side. Taking into account that unitarity fixes to unity the perimeter of the triangle, we conclude that the values of the moduli for which cancellation is possible must satisfy:

$$\begin{aligned} \frac{1}{2} &\geq |U_{e1}|^2, \\ \frac{1}{2} &\geq |U_{e2}|^2 \geq \frac{1}{2} - |U_{e1}|^2. \end{aligned} \quad (13)$$

Assuming instead that a signal is observed, the upper bound on the elements of the mixing matrix resulting from geometrical considerations reads:

$$|U_{ej}|^2 \leq \frac{1 + |\mathcal{M}_{ee}|/m_0}{2}. \quad (14)$$

Even with a maximal value of the mass ratio:  $|\mathcal{M}_{ee}|/m_0 = 30 \%$  (that would imply that a signal of  $0\nu\beta\beta$  will be observed close to the present limit) the upper bound does not change much:  $|U_{ej}|^2 \leq 0.65$ .

---

<sup>3</sup>In last analysis these parameters have be related to instrumental or maybe new physics effects.

We want to make one point clear: we are not proposing any explanations of the cancellation between the elements of the mixing matrix squared, we are just remarking that, once the hypotheses underlined above are accepted, consistency with the data *requires a cancellation*; this is true even if a positive signal, related to a small mass  $\mathcal{M}_{ee}$  should be found. We show below that quite a precise pattern of the mixing matrix can be figured out insisting on this scheme, and on the consistency with the experimental data.

Eq. (13) implies that there is at most one small mixing matrix element  $U_{ej}$ , and even in this case the other two elements correspond to maximal mixing angle. The presence of large mixings, that was generally noticed in previous studies [1–12] is simply a consequence of the absence of signal in  $0\nu\beta\beta$  searches in these models.

It was remarked that the scenarios with large lepton mixing tend to be disfavoured by the comparison of the observed SN1987A neutrino energy spectrum and the theoretical spectra [38]. Therefore, a clear understanding of the mechanism of neutrinos emission and propagation in the process of stellar collapse could provide us with the most direct way to exclude the neutrino scenario under discussion. We will not speculate further on this idea in the present work, and concentrate instead on the study of possible patterns of the mixing matrix.

### C. Information from the reactors

The neutrino beams produced at nuclear reactors are well suited for the study of disappearance of electron neutrino flux. The neutrino oscillations are described by the conversion probability

$$P_{ee} = 1 - 4 \sum_{j>i} |U_{ej}|^2 |U_{ei}|^2 \sin^2 \left[ \frac{\Delta m_{ji}^2 L}{4 p} \right], \quad (15)$$

which just depend on the mixing angles which are constrained by the neutrinoless double beta decay. In the case A, since that the parameter  $\Delta m_{\odot}^2$  is too small to develop oscillating phases on terrestrial distances, our expression (15) reduces to:

$$P_{ee} = 1 - 4 |U_{e3}|^2 (1 - |U_{e3}|^2) \sin^2 \left[ \frac{\Delta m_{atm}^2 L}{4 p} \right] \quad (\text{case A}) \quad (16)$$

that is effectively a two-flavour neutrino task, the relevant element of the mixing matrix being the one related with the “isolated” massive neutrino (in case B, one arrive at the same expression with  $U_{e3}$  replaced by  $U_{e1}$ ). The neutrino flux is not suppressed at all in the two extreme cases:  $|U_{e3}|^2 \approx 0$  or  $\approx 1$ , but the second solution must be discarded in view of the absence of  $0\nu\beta\beta$  signal, that implies eq. (14). Applying the inequalities (13) we conclude:

$$|U_{e1}|^2 \approx 1/2, \quad |U_{e2}|^2 \approx 1/2, \quad |U_{e3}|^2 \approx 0 \quad (\text{case A}) \quad (17)$$

(in case B the small element is  $|U_{e1}|$ ). The constraints from reactor experiments are not, of course, absolute. To be quantitative we notice that for a value of  $\Delta m_{atm}^2 = 0.01 \text{ eV}^2$  that may explain well the atmospheric anomaly we get the bound [39]:

$$|U_{e3}|^2 < 0.2 \text{ at } 90\% \text{ CL} \quad (\text{case A}). \quad (18)$$

This bound however depends strongly on the value of  $\Delta m_{atm}^2$ , and becomes already ineffective if instead of 0.01 we have  $7.5 \times 10^{-3} \text{ eV}^2$ : This should warn from a literal use of equation (17) (compare also with the discussion in [12]). The future search at Palo Verde [40] and at CHOOZ [41]) will be able to strengthen the bound, or maybe suggest a non-zero value of the mixing angle. Assuming eq. (17), the bounds from neutrinoless double beta decay are easily solved:  $U_{e2} \approx \pm i \times U_{e1}$ , where  $i$  is the imaginary unity.

### III. FEATURES OF THE SCENARIO

Let us summarize the major consequences of the constraints exploited in previous section by using the matrix  $|\mathcal{U}|$ , whose elements are the numbers  $|U_{ej}|$ . Using eq. (17), assuming for simplicity that  $U_{e3}$  is quite small—our stronger hypothesis—, and the unitarity relations<sup>4</sup>, the approximate pattern is:

---

<sup>4</sup>In strict analogy with the what happens to the Cabibbo-Kobayashi-Maskawa mixing matrix, the matrix  $|\mathcal{U}|$  can be described in the general case by four parameters (for instance one can use three mixing angles and one phase).

$$|\mathcal{U}| \approx \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{\sin \theta}{\sqrt{2}} & \frac{\sin \theta}{\sqrt{2}} & \cos \theta \\ \frac{\cos \theta}{\sqrt{2}} & \frac{\cos \theta}{\sqrt{2}} & \sin \theta \end{pmatrix} \quad (\text{case A}) \quad (19)$$

(in case B, the last column exchanges with the first one). There is only one parameter in eq. (19),  $\theta \in [0, \pi/2]$ ; we will see in the following discussion that the preferred value is  $\theta \approx \pi/4$ . As discussed in sub-section C. above, the zero entry should be intended as an approximate statement only. In the rest of this work we will discuss the qualitative features of the scenario outlined, leaving the detailed comparison with the data for a future work.

### A. Atmospheric neutrinos

On terrestrial distances  $L \approx R_e$ , and for neutrino momenta  $p \approx 1$  GeV the oscillations related to the mass difference  $\Delta m_\odot$  do not develop:  $\cos(\Delta m_\odot^2 L/(2p)) \rightarrow 1$ . Therefore, in case A, the expressions for the matrix  $\mathcal{P}$  of the conversion probabilities (whose elements are  $P_{\ell\ell'}$ , where  $\ell$  and  $\ell' = e, \mu, \tau$ ) is:

$$\mathcal{P} \approx \mathbb{1} - \sin^2 2\theta \times \sin^2 \left[ \frac{\Delta m_{atm}^2 L}{4p} \right] \times \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix}. \quad (20)$$

If we suppose that the oscillations related to  $\Delta m_{atm}^2$  are effectively averaged, at least at distances  $L$  of the order of the Earth size, we can replace the oscillating factor  $\sin^2(\Delta m_{atm}^2 L/(4p))$  by  $1/2$ . A rough estimation of the muon-to-electron neutrino ratio  $r = \Phi(\nu_\mu)/\Phi(\nu_e)$  measured in the detectors can be obtained assuming that in the production region  $r_0 = \Phi_0(\nu_\mu)/\Phi_0(\nu_e) \approx 2$ , that is adequate for neutrinos of energies around 1 GeV originated by cosmic rays entering the Earth atmosphere. Using  $\Phi(\nu_\ell) = \sum_{\ell'=e,\mu,\tau} P_{\ell\ell'} \Phi_0(\nu_{\ell'})$  we find for the double-ratio  $R$ :

$$R \equiv \frac{r}{r_0} \approx 1 - \frac{\sin^2(2\theta)}{2}. \quad (21)$$

Kamiokande [42] and Superkamiokande [43] measurements in the multi-GeV energy region suggest  $R \approx 0.55$ , and this implies that  $\theta$  has to be close to  $\theta = \pi/4$  to produce a large suppression factor. However, the data are much better fitted if a linear dependence of the spectrum from the cosine of the zenith angle  $\phi_Z$  is also introduced:

$$R(\phi_Z) \approx 0.55 + 0.3 \times \cos \phi_Z. \quad (22)$$

This favours not-so-large values of  $\Delta m_{atm}^2$ , in such a way that the averaging effect is reduced for smaller neutrino travel distance (around 10 – 100 km, corresponding to  $\phi_Z$  in a neighbourhood of 0) and it is maximal for neutrinos produced in the antipodal regions (produced at  $L \approx 10000$  km far apart from the detector, that is for  $\phi_Z \approx \pi$ ). So the averaged oscillations would take place at maximal distances only:  $R(\pi) \approx 0.3$ , which should be compared with the smallest possible theoretical value,  $R \approx 0.5$ . This is not excluded by the data, the errors in each of the  $5 \times 2$  data points being around 0.1 (except for  $\phi_Z \approx 0$ , where the errors are 4 – 5 times larger). But while waiting for more precise data, one has to keep in mind that the model considered suggests somewhat smaller flux suppression than measured at present.

### B. Solar neutrinos

We consider the vacuum oscillation mechanism as a solution of the solar neutrino problem, which requires  $\Delta m_\odot^2 \approx 10^{-10}$  eV<sup>2</sup>. The survival probability of electron neutrinos in case A is:

$$P_{ee} = 1 - 2|U_{e3}|^2(1 - |U_{e3}|^2) - 4|U_{e1}|^2|U_{e2}|^2 \sin^2 \left[ \frac{\Delta m_\odot^2 L}{4p} \right] \quad (23)$$

where  $L$  represents the Earth-Sun distance. The second term gives the oscillating factor, whereas the first term can give an additional (constant in energy) suppression. In fact, if  $|U_{e1}|^2 = |U_{e2}|^2$  we can rewrite:

$$P_{ee} = |U_{e3}|^4 + (1 - |U_{e3}|^2)^2 \cos^2 \left[ \frac{\Delta m_\odot^2 L}{4p} \right]. \quad (24)$$

If the bound (18) holds, the first term is negligible; therefore the oscillating factor can be multiplied by a suppression factor that can be as small as 0.64. This can somewhat diminish the spectrum distortion and the seasonal variations that are correlated characteristics of the vacuum-oscillation mechanism [45]. If, beside that, we suppose that the oscillations are effectively averaged (requiring that the splitting  $\Delta m_\odot^2$  is large in comparison with  $10^{-10}$  eV<sup>2</sup>), the probability of suppression is close to the maximal *averaged* value  $\langle P_{ee} \rangle_{max} = 1/3$ . This may be of relevance for neutrinos emitted by a supernova, or maybe for solar neutrinos themselves [46].

#### IV. SUMMARY AND DISCUSSION

We summarize our findings in fig. 1, which visualizes the flavour content  $|U_{\ell i}|^2$  of the massive neutrinos  $\nu_i$  in the way used in [44] to present a panorama of scenarios of neutrino masses and mixing.




FIG. 1. Composition of massive neutrinos: white for electron-, hatched for muon- and black for tau-neutrino flavour. For a proper interpretation of the vertical scale, recall that the mass splittings, eqs. (7) are much smaller than the common mass scale  $m_0$ , eq. (9).

Let us recall the main features of the scenario proposed, that assumes that the solar and the atmospheric anomalies in the fluxes of neutrinos are due to oscillations:

1. Cosmological indications can be reconciled with the the solar and atmospheric neutrino deficits if the three Standard Model neutrinos have in first approximation the same mass  $m_0 = m(\text{cosm.})$ .
2. We emphasized that the common mass  $m_0$  needed in hot+cold DM scenarios can be probed in direct searches for neutrino masses, and that this mass is only *two times smaller* than the masses that can be currently excluded (eqs. (9) and (11)). The case in which the energy of the Universe is dominated by three massive neutrinos can already be excluded.
3. The absence of a signal in neutrinoless double beta decay searches requires a cancellation. Even if this is not *a priori* a theoretical attractive perspective, it implies strong restrictions on certain combinations of the elements of the mixing matrix (compare with eq. (13)).
4. The absence of signal in reactor experiments suggests that the structure of the mixing matrix, eq. (19), may be quite simple (see also eq. (18) and related discussion). Regarding this point, future electron neutrino disappearance experiments will be important to better fix the parameters of the model. Even better, they could permit us to distinguish the present variant of the scenario with nearly degenerate neutrinos with the variant in which the solar neutrino deficit is explained invoking the MSW effect; in fact, in this latter case *large* values of the mixing element  $U_{e3}$  are preferred [12] (compare with eq. (16) and discussion thereafter).
5. The pattern of the mixing matrix allows us to take into account the atmospheric neutrino anomaly. With accumulation of data at SuperKamiokande, the zenith angle dependence of the neutrino spectrum will become a severe test of the scenario proposed.
6. Regarding the observed suppression of the solar neutrinos flux, we remarked that this model can incorporate the vacuum oscillation solution, possibly with reduced distortion of the energy spectrum.

In conclusion, we express our opinion that, even if the model proposed is not at present of particular theoretical appeal, it is of some interest since (a) it incorporates cosmological and phenomenological indications; (b) it is enough restrictive to be confirmed or disproved in the future (few) years.

## ACKNOWLEDGMENTS

I am grateful to the Theory Group of FERMILAB, and in particular to the Deputy Head of the High-Energy section R.K. Ellis, for hospitality during the last stages of this work. It is also a pleasure to thank J. Peltoniemi (who created and maintains the database *The Ultimate Neutrino Page* at the URL <http://www.physics.helsinki.fi/neutrino/>) for scientific help and also for encouragement; R.N. Mohapatra, M. Persic, A. Riotto, E. Roulet, P. Salucci and A. Stebbins for discussions about aspects of cosmology; E.Kh. Akhmedov and M.V. Chizhov for discussions about neutrino phenomenology. I need special thanks for A.Yu. Sminorv, who introduced me to the subject of neutrino physics and helped me to orient in this vast field.

- 
- [1] D.O. Caldwell and R.N. Mohapatra, Phys.Rev. D48 (1993) 3259.
  - [2] S.T. Petcov and A.Yu. Smirnov, Phys.Lett. B322 (1994) 109.
  - [3] A.S. Joshipura, Zeit.Phys. C64 (1994) 31.
  - [4] D.O. Caldwell and R.N. Mohapatra, Phys.Rev. D50 (1994) 3477.
  - [5] A. Ioannisian and J.W.F. Valle, Phys.Lett. B332 (1994) 93.
  - [6] B. Bamert and C.P. Burgess, Phys.Lett. 329 (1994) 289.
  - [7] D.G. Lee and R.N. Mohapatra, Phys.Lett. B329 (1994) 463.
  - [8] A.S. Joshipura, Phys.Rev. D51 (1995) 1321.
  - [9] S. Nussinov and R.N. Mohapatra, Phys.Lett. B346 (1995) 75.
  - [10] K.S. Babu, R.K. Schaefer and Q. Shafi, Phys.Rev. D53 (1996) 606.
  - [11] P. Binétruy, S. Lavignac, S.T. Petcov and P. Ramond, Nucl.Phys. B496 (1997) 3.
  - [12] H. Minakata and O. Yasuda, Phys.Rev. D56 (1997) 1692.
  - [13] S.P. Mikheyev and A.Yu. Smirnov, Yad.Fiz., 42 (1985) 1441.
  - [14] S.P. Mikheyev and A.Yu. Smirnov, Nuovo Cim. 9C (1986) 17.
  - [15] L. Wolfenstein, Phys.Rev. D17 (1978) 2369.
  - [16] B. Pontecorvo, Zh.Exp.Teor.Fiz. 53 (1967) 1717.
  - [17] Review on *Dark Matter* by M. Srednicki in [18] (page 116).
  - [18] R.M. Barnett et al. (Particle Data Group collaboration), Phys.Rev. D 54 (1996) 1.
  - [19] E.L. Wright et al. (COBE collaboration), Ap.J. 396 (1992) L13, and the references to the theoretical models quoted therein.
  - [20] J.R. Primack, J. Holtzman, A. Klypin and D.O. Caldwell, Phys.Rev.Lett. 74 (1995) 2160. This work suggests the scenario in which the hot DM is formed by two massive neutrinos, equally sharing 5 eV of total mass.
  - [21] Review on *The Hubble Constant* by C.J. Hogan in [18] (page 112).
  - [22] W.L. Freedman, *Determination of the Hubble Constant* astro-ph/9612024, presented at the Conference Critical Dialogs in Cosmology, June 1996, Princeton, NJ.
  - [23] J.R. Primack, *Dark Matter and Structure Formation*, astro-ph/9707285, presented at the Jerusalem Winter School 1996.
  - [24] S. Weinberg, *Gravitation and Cosmology*, John Wiley & sons press (1972).
  - [25] A. Albrecht and A. Stebbins, Phys.Rev.Lett. 69 (1992) 2615. However, recent numerical simulations [26] suggest that cosmic strings may not be able to account for the structure at large scales, and this could require major modifications of this scenario.
  - [26] U.-L. Pen, U. Seljak and N. Turok, astro-ph/9704165.
  - [27] G. Raffelt and J. Silk, Phys.Lett. B192 (1987) 65.
  - [28] J. Peltoniemi and J.W.F. Valle, Nucl.Phys. B406 (1993) 409.
  - [29] J. Peltoniemi, D. Tommasini and J.W.F. Valle, Phys.Lett. B298 (1993) 383.
  - [30] J. Peltoniemi, Mod.Phys.Lett. A8 (1993) 3593.
  - [31] C. Athanassopoulos et al. (LSND collaboration), Phys.Rev.Lett. 77 (1996) 3082.
  - [32] C. Athanassopoulos et al. (LSND collaboration), Phys.Rev. C54 (1996) 2685.
  - [33] C. Athanassopoulos et al. (LSND collaboration), nucl-ex/9706006.
  - [34] K. Eitel (for KARMEN collaboration), hep-ex/9706023, presented at the 32<sup>nd</sup> Rencontres de Moriond: Electroweak Interactions and Unified Theories, March 1997, Les Arcs, France.



- [35] V.M. Lobashev (Troitsk collaboration), preliminary data analysis presented at the XVI International Workshop on Weak Interactions and Neutrinos, 22 – 29 June 1997, Capri, Italy [36].
- [36] Copies of the transparencies can be found at the link <http://www.na.infn.it/win97/>.
- [37] A. Balysh et al. (Heidelberg-Moskow experiment), Phys.Lett. B356 (1995) 450.
- [38] A.Yu. Smirnov, D.N. Spergel and J.N. Bahcall, Phys.Rev. D49 (1994) 1389.
- [39] G.S. Vidyakin et al. (Krasnoyarsk collaboration) JETP Lett. 59 (1994) 390.
- [40] F. Bohm et al. (Palo Verde collaboration), STANFORD-HEP-96-04, presented at the 17<sup>th</sup> International Conference on Neutrino Physics and Astrophysics, June 1996, Helsinki, Finland.
- [41] R.I. Steiberg, hep-ph/9306282, presented at the 5<sup>th</sup> International Workshop on Neutrino Telescopes, March 1993, Venice, Italy.
- [42] Y. Fukuda et al. (Kamiokande collaboration), Phys.Lett. B335 (1994) 237.
- [43] K. Nakamura (SuperKamiokande collaboration), preliminary data analysis presented at the XVI International Workshop on Weak Interactions and Neutrinos, 22 – 29 June 1997, Capri, Italy [36].
- [44] A.Yu. Smirnov, *Neutrino Masses and Mixing*, hep-ph/9611465, presented at the 28<sup>th</sup> International Conference on High-Energy Physics (ICHEP 96), Warsaw, Poland.
- [45] S.P. Mikheyev and A.Yu. Smirnov, hep-ph/9708403.
- [46] A. Acker and S. Pakvasa, Phys.Lett. B397 (1997) 209.